## Patent claims:

5

10

15

- A process for the storage of a protein in an aqueous solution, comprising the addition of cysteine to the aqueous solution, wherein said addition results in a delay in the temporal decrease in the effective concentration of the protein.
  - The process as claimed in claim 1, wherein the protein is a heterologous protein prepared in a microorganism.
  - The process as claimed in claim 2, wherein the microorganism is a bacterium.
- The process as claimed in claim 3, wherein the bacterium is Escherichia coli.
  - The process as claimed in claim 2, wherein the microorganism is a yeast.
- The process as claimed in claim 5, wherein the yeast is Saccharomyces cerevisiae.
  - 7. The process as claimed in claim 5, wherein the yeast is Pichia pastoris.
- 25 8. The process as claimed in claim 1, wherein the protein is a heterologous protein and is prepared in an insect cell.
  - The process as claimed in claim 2 or claim 8, wherein the protein is prepared from an expression vector construct.

10

20

- The process as claimed in claim 1, wherein the protein is present in dissolved form.
- The process as claimed in claim 1, wherein the protein is present in suspension.
  - The process as claimed in claim 1, wherein the concentration of cysteine in the aqueous protein solution is in the range from about 100 mM to about 500 mM.
  - The process as claimed in claim 12, wherein the concentration of cysteine in the aqueous protein solution is in the range from about 150 mM to about 220 mM.
- 15 14. The process as claimed in claim 12, wherein the concentration of cysteine in the aqueous protein solution is about 170 mM.
  - 15. The process as claimed in claim 1, wherein the storage of the protein takes place at about 0°C to about 50°C.
  - The process as claimed in claim 15, wherein the storage of the protein takes place at about 5°C to about 30°C.
- 17. The process as claimed in claim 16, wherein the storage of the protein25 takes place at about 5°C.
  - The process as claimed in claim 1, wherein the protein stored is insulin, an insulin derivative, or a precursor thereof.

5

10

15

- 19. A process for the preparation and storage of a heterologous protein, comprising the expression of the heterologous protein or its precursor in a transformed microorganism, optional disruption of the microorganism and/or isolation of the heterologous protein or its precursor from the culture medium, and the subsequent storage of the heterologous protein according to the process of claim 1.
- 20. The process of claim 19, further comprising the renaturation of the heterologous protein or its precursor and the purification and isolation of the heterologous protein, including optional removal of a leader sequence or other sequences that may be present in the precursor of the heterologous protein.
- The process as claimed in claim 19, wherein the heterologous protein is animal insulin.
  - The process as claimed in claim 20, wherein the animal insulin is human insulin.